Using Map of Dermatology (http://www.healthcybermap.org/dermap/) answer the following MCQs:

(Level: easy; undergraduate students and non-specialist clinicians)

- 1. Erythema nodosum typically occurs on:
- a. Face
- b. Trunkc. Hands
- d. Legs
- 2. Tuberculosis verrucosa cutis occurs most commonly on:
- b. Neck
- c. Trunk
- d. Hands
- 3. The following conditions may present as vegetative lesions on the hands, except:
- a. Erysipeloid
- b. Keratoacanthoma
- c. Squamous cell carcinoma
 d. Verruca vulgaris
- 4. Which of the following conditions does not affect the palms:
- a. Dyshidrotic eczematous dermatitis
- b. Erythema multiforme
- c. Pustular psoriasis
- d. Seborrheic keratosis
- 5. Which of the following oculomucocutaneous syndromes is sometimes associated with Keratoderma blennorrhagicum on the soles of the feet:
- a. Behçet's disease
- b. Erythema multiforme
- c. Reiter's syndromed. Sweet's syndrome
- 6. Discrete inflammatory-type yellow papules appearing suddenly on the buttocks can be a presentation of:
 - a. Acne
- b. Eruptive xanthoma
- c. Folliculitis
- d. Scabies
- 7. An obese diabetic patient is presenting with diffuse velvety thickening and hyperpigmentation in the axillae. The most likely diagnosis is:
- a. Acanthosis nigricans
- b. Erythrasmac. Hidradenitis suppurativa
- d. Pityriasis versicolor
- 8. Macula cerulea is seen in:
- a. Pediculosis capitis
- b. Pediculosis pubis
- c. Meningococcemia
- d. Minocycline pigmentation (ceruloderma)
- 9. Erythema migrans on the trunk is a feature of:
- a. Drug eruption
- b. Lyme borreliosis
- c. Mastocytosis
- d. Tinea corporis
- 10. Depigmented ash-leaf macules on the trunk are often the earliest cutaneous feature of:
- a. LEOPARD syndrome
- b. Leprosy
- c. Neurofibromatosis
- d. Tuberous sclerosis

1(d), 2(d), 3(a), 4(d), 5(c), 6(b), 7(a), 8(b), 9(b), 10(d)